

Exact results for spatial decay of correlations in low-dimensional insulators II

Janusz Jędrzejewski^{1*} and Taras Krokhmal'skii^{2,1†}

¹*Institute of Theoretical Physics, University of Wrocław, pl. Maksa Borna 9, 50-204 Wrocław, Poland*

²*Institute for Condensed Matter Physics, 1 Svientsitskii Str., L'viv-11, 79011, Ukraine*

(Dated: February 2, 2008)

We study decay rates of one-body reduced density matrices in insulators, described by a tight-binding model, where not only an external potential but also hoppings are spatially modulated. We determine analytically the power in the power law and the correlation length in $D = 1$ case and in several lattice directions in $D = 2$ case. Unlike the uniform hopping case, in $D = 1$ system and in some directions of $D = 2$ system the correlation length is not determined uniquely by the gap. Moreover, a crossover from $D = 2$ -decay rates to $D = 1$ ones is investigated.

PACS numbers: 71.10.Fd, 71.20.-b

The localization properties of one-body reduced density matrices (DM) are of vital importance for rapidly developing computational techniques, used for calculating various characteristics of crystalline solids^{1,2,3}. But the interest in large-distance decay rates of DM dates from 1959, when Kohn⁴ determined an exponential factor in the large-distance asymptotics of DM in one-dimensional ($D = 1$) crystals. Recent works concerning this problem include Refs.^{5,6,7,8,9,10,11,12}. In particular, another major step, after Kohn's result, was made by He and Vanderbilt who discovered a power-law factor in the decay rate of $D = 1$ crystals. Extensions to systems in dimensions $D > 1$ have been obtained by Des Cloizeaux¹³, Taraskin et al^{10,11}, and in our recent paper¹², where we have shown to what extent the large-distance asymptotics of DM is sensitive to lattice directions in $D = 2$ crystals.

In all the papers quoted above, the pattern of hoppings is spatially uniform. Here, we consider a system of spinless fermions on a lattice whose hoppings are spatially modulated, subjected to a modulated external potential. The Hamiltonian of the considered system reads:

$$H = \sum_i U_i a_i^\dagger a_i + \sum_{\langle i, j \rangle} t_{ij} (a_i^\dagger a_j + h.c.). \quad (1)$$

In the above expression, i, j represent the lattice sites of a D -dimensional simple cubic lattice, $\langle i, j \rangle$ stand for the pairs of nearest neighbors on this lattice, and in the sum over nearest neighbors each pair is counted once. The operators $a_i^\dagger$ (a_i) create (annihilate) a spinless fermion in a single-particle orbital $|i\rangle$, belonging to an orthonormal basis. As j runs through the underlying lattice, the value of the external potential at site j is modulated with the amplitude u ,

$$U_j = U + (-1)^{\sigma_j} u, \quad (2)$$

where for a site $j = (j_1, \dots, j_D)$, $\sigma_j = \sum_{l=1}^D |j_l|$ is the (noneuclidean) distance of this site from the origin. Since a D -dimensional simple cubic lattice consists of two interpenetrating sublattices: the even one – with σ_j even, and the odd one – with σ_j odd, the external potential is uniform at each sublattice. Its value changes by $\pm 2u$ on passing from one sublattice to the other one, and the

average over the whole lattice is U . Similarly, the symmetric hopping matrix t_{ij} is modulated:

$$t_{ii \pm e_l} = t [1 \pm (-1)^{\sigma_i} \tau], \quad (3)$$

with the mean t and the amplitude $t\tau$, where e_l , $l = 1, \dots, D$, is the unit vector with only one nonzero component – the l -th one. Without any loss of generality, we can suppose that $u \geq 0$, $t \geq 0$, and $1 \geq \tau \geq 0$.

The considered model can describe electronic properties of crystal surfaces, possibly containing adsorbed atoms. The pattern of hoppings (3) may arise when atoms form a distorted square lattice, that is one of the sublattices is shifted with respect to the other.

Under the periodic boundary conditions, the Hamiltonian (1) is block-diagonalized by the plane wave orbitals, $|\mathbf{k}\rangle$, with the wave vector in the first Brillouin zone of the lattice. Specifically, shifting zero of the energy scale to U and expressing all the energies involved in the units of the average transfer integral t , we obtain the upper, $\Lambda_{\mathbf{k}}^+$, and the lower, $\Lambda_{\mathbf{k}}^-$, bands of eigenenergies:

$$\Lambda_{\mathbf{k}}^\pm = \pm \sqrt{u^2 + \varepsilon_{\mathbf{k}}^2 + \zeta_{\mathbf{k}}^2} \equiv \pm \Delta_{\mathbf{k}}. \quad (4)$$

In Eq.(4), the wave vector $\mathbf{k}$ is restricted to the first Brillouin zone of one of the sublattices (so that there is as many eigenvalues as sites in the lattice), $\varepsilon_{\mathbf{k}} = 2 \sum_{l=1}^D \cos k_l$, $\zeta_{\mathbf{k}} = 2\tau \sum_{l=1}^D \sin k_l$. The two bands are mutually symmetric about zero and are separated by a gap g_D . If the system given by (1) is half-filled and $g_D > 0$, then it is an insulator in the sense of the standard band theory. The normalized eigenvectors corresponding to the eigenvalues $\Lambda_{\mathbf{k}}^\pm$ are linear combinations of the vectors $|\mathbf{k}\rangle$ and $|\mathbf{k} + \boldsymbol{\pi}\rangle$, where $\boldsymbol{\pi}$ stands for the vector with all the components equal to π . By means of these eigenvectors, one can calculate the zero-temperature non-diagonal elements of DM for the half-filled system, which in the thermodynamic limit read

$$\begin{aligned} \langle a_i^\dagger a_{i+\mathbf{r}} \rangle &= -\frac{1}{2} \sum_{l=1}^D [\mathcal{S}_l \mathcal{R}_D(\mathbf{r}) - \tau (-1)^{\sigma_i} \mathcal{D}_l \mathcal{R}_D(\mathbf{r})], \\ &\quad \text{if } \sigma_{\mathbf{r}} = 2m+1, \text{ and} \\ \langle a_i^\dagger a_{i+\mathbf{r}} \rangle &= -\frac{u}{2} (-1)^{\sigma_i} \mathcal{R}_D(\mathbf{r}), \text{ if } \sigma_{\mathbf{r}} = 2m, \end{aligned} \quad (5)$$

where

$$\mathcal{R}_D(\mathbf{r}) = 2(2\pi)^{-D} \int_{B.Z.} d\mathbf{k} \exp(i\mathbf{k}\mathbf{r}) \Delta_{\mathbf{k}}^{-1}, \quad (6)$$

and $\mathcal{S}_1\mathcal{R}(\mathbf{r}) \equiv \mathcal{R}(r_1 + 1, \dots, r_D) + \mathcal{R}(r_1 - 1, \dots, r_D)$, $\mathcal{D}_1\mathcal{R}(\mathbf{r}) \equiv \mathcal{R}(r_1 + 1, \dots, r_D) - \mathcal{R}(r_1 - 1, \dots, r_D)$, and so on for $l = 2, \dots, D$. In Eq. (6), the D-dimensional integral is taken over the first Brillouin zone of a sublattice and $\mathbf{r} \neq 0$. At odd sites ($\sigma_{\mathbf{r}} = 2m + 1$), the function $\mathcal{R}_D$ vanishes.

At the level of the Hamiltonian (1), it is clear that for $\tau = 0$ we are back in the the uniform hopping case of Ref.¹². Note however, that here $\Delta_{\mathbf{k}}$ is twice as large as in Ref.¹², and consequently an extra factor 2 appears in the definition of $\mathcal{R}_D(\mathbf{r})$. On the other hand, if $\tau = 1$, the system loses its connectivity with respect to the hopping. Specifically, for $D = 1$, we obtain a system of independent dimers whose correlation functions vanish at distances greater than one, while for $D = 2$ – a system of independent quasi-one-dimensional zig-zag chains.

In what follows, we shall investigate the large-distance decay rate of the function $\mathcal{R}_D(\mathbf{r})$ in $D = 1, 2$ systems.

It is instructive to analyze first the one-dimensional case. The gap in the spectrum is $g_1 = 4\delta_1$, with $\delta_1^2 = \beta_1^2 + \tau^2$, and $\beta_1 = u/2$. Apparently, the role of the rescaled modulation amplitudes, u and τ , in creating the gap is symmetric. Moreover, $g_1 > 0$ unless $u = \tau = 0$. For later purposes, it is convenient to introduce, besides the rescaled gap δ_1 , one more parameter describing the spectrum: for the lower band this parameter amounts to the ratio, κ_1 , of the upper edge of the band to the lower edge, i.e. $\kappa_1^2 = (1 - \tau^2)/(1 + \beta_1^2)$. The pair (δ_1, κ_1) can be thought of as new coordinates, equivalent to the coordinates (β_1, τ) . The specific combination of β_1 and τ , denoted κ_1 , appears naturally in an expression for the correlation function (5). Provided that $\kappa_1 \neq 0$, the function $\mathcal{R}_1(\mathbf{r})$ at even points, $\mathbf{r} = 2m$, can be written as

$$\begin{aligned} \mathcal{R}_1(2m) &= \frac{1}{2\pi} \frac{1}{\sqrt{1 + \beta_1^2}} \int_{-\pi/2}^{\pi/2} dk \frac{\cos(2mk)}{\sqrt{1 - \kappa_1^2 \sin^2 k}} \\ &= \frac{(-1)^m}{\pi} \frac{1}{\sqrt{1 - \tau^2}} Q_{m-1/2} \left(\frac{2}{\kappa_1^2} - 1 \right), \end{aligned} \quad (7)$$

where Γ stands for the Euler Γ -function, F – for the Gauss hypergeometric function, and Q_ν – for the Legendre function of second kind¹⁴. Eq.(7) implies the following large- m asymptotics:

$$\mathcal{R}_1(2m) \approx \frac{(-1)^m}{\sqrt{2\pi}} \frac{(1 - \kappa_1^2)^{1/4}}{\delta_1} \frac{\exp(-2m/\xi)}{\sqrt{2m}}, \quad (8)$$

where the inverse correlation length ξ^{-1} is given by

$$\xi^{-1} \equiv \ln \frac{1 + \sqrt{1 - \kappa_1^2}}{\kappa_1}. \quad (9)$$

We note that the correlation function $\langle a_i^\dagger a_{i+r} \rangle$ is continuous at $\tau = 0$, $\tau = 1$, and $\tau = \beta_1 = 0$. Out of the

three above cases, the asymptotics (8) is continuous only at $\tau = 0$, where we recover the asymptotics obtained in Ref.¹².

The new variables (δ_1, κ_1) reveal clearly the nature of the relation between ξ , κ_1 and δ_1 . It follows from (9) that κ_1 determines uniquely ξ and vice versa. But the relation between ξ (or κ_1) and δ_1 is a multi-valued one. Specifically, to given δ_1 there corresponds an interval $[\kappa_1^{min}(\delta_1), \kappa_1^{max}(\delta_1)]$ of κ_1 values, where $\kappa_1^{min}(\delta_1) = \sqrt{1 - \delta_1^2}$ if $\delta_1 \leq 1$, and $\kappa_1^{min}(\delta_1) = 0$ if $\delta_1 > 1$, and $\kappa_1^{max}(\delta_1) = 1/\sqrt{1 + \delta_1^2}$. Consequently, the inverse correlation length ξ^{-1} can assume any value from the interval $[\xi^{-1}(\kappa_1^{max}(\delta_1)), \xi^{-1}(\kappa_1^{min}(\delta_1))]$. As δ_1 tends to zero, the edges of the interval tend to zero as well but its width shrinks even faster, what results in a fairly good localization of ξ^{-1} , for small δ_1 . Specifically, expanding the bounds at $\delta_1 = 0$,

$$\xi^{-1}(\kappa_1^{max}(\delta_1)) \approx \delta_1 - \frac{1}{6}\delta_1^3 + \frac{3}{40}\delta_1^5 + \dots, \quad (10)$$

$$\xi^{-1}(\kappa_1^{min}(\delta_1)) \approx \delta_1 + \frac{\delta_1^3}{3} + \frac{\delta_1^5}{5} + \dots, \quad (11)$$

we see that to the first order in δ_1 the bounds coincide and the indeterminacy-interval width is $\delta_1^3/2$, in the leading order.

For $\delta_1 > 1$, ξ^{-1} can assume any value not less than $\xi^{-1}(\kappa_1^{max}(\delta_1))$, and this lower bound blows up as $\delta_1 \rightarrow \infty$.

The described above indeterminacy of ξ^{-1} vanishes if one of the parameters, β_1 or τ , is kept constant, since such conditions turn the relations between κ_1 and δ_1 into single valued functions. Of particular interest is the behavior of ξ^{-1} in two such situations, motivated physically. Firstly, a weak modulated external potential (β_1 varies in a vicinity of zero) is imposed on a connected system with modulated hopping ($0 < \tau < 1$, fixed). Secondly, a weak modulated hopping is imposed (τ varies in a vicinity of zero) on a system in a modulated potential ($0 < \beta_1$, fixed). In the both cases, δ_1 is separated from zero. Therefore, to describe the behavior of ξ^{-1} it is more convenient to consider the expansions in β_1 or τ , respectively. Specifically, in the first situation

$$\xi^{-1} \approx \ln \sqrt{\frac{1 + \tau}{1 - \tau}} + \frac{1}{\tau} \frac{\beta_1^2}{2} - \frac{1 + \tau^2}{\tau^3} \frac{\beta_1^4}{8} + \dots, \quad (12)$$

while in the second one

$$\xi^{-1} \approx \ln(\beta_1 + \sqrt{1 + \beta_1^2}) + \frac{\sqrt{1 + \beta_1^2}}{\beta_1} \frac{\tau^2}{2} + \dots \quad (13)$$

Clearly, in the above expansions it is not possible to go to zero with the fixed parameters. To obtain the corresponding expansions about zero, we note that the function $\kappa_1^{max}(\delta_1) = \kappa_1(\beta_1, \tau)|_{\beta_1=\delta_1, \tau=0}$. Therefore, for $\tau = 0$ the expansion in β_1 coincides with the expansion (10) of the lower bound for ξ^{-1} . Similarly, $\kappa_1^{min}(\delta_1) =$

$\kappa_1(\beta_1, \tau)|_{\beta_1=0, \tau=\delta_1}$, hence for $\beta_1 = 0$ the expansion in τ coincides with the expansion (11) of the upper bound for ξ^{-1} .

Yet another interesting regime is, when a connected system ($\tau \neq 1$) is subjected to a strongly modulated external potential. Then, the large β_1 asymptotics of ξ^{-1} reads:

$$\xi^{-1} \approx \ln \frac{2}{\sqrt{1-\tau^2}} + \ln \beta_1 + \frac{1+\tau^2}{4\beta_1^2} + \dots \quad (14)$$

The two-dimensional system differs from the one-dimensional one in many respects, due to different geometry and connectivity of the underlying lattice. When hopping is modulated, an important difference appears already in the spectrum: the gap $g_2 = 8\beta_2$, $\beta_2 = u/4$, is independent of the amplitude τ , and vanishes unless the external potential is modulated.

The function $\mathcal{R}_2(\mathbf{r})$ can be cast into the form

$$\mathcal{R}_2(\mathbf{r}) = \frac{1}{\pi^2} \int_0^{\frac{\pi}{2}} \int_0^{\frac{\pi}{2}} \frac{dx dy \cos(2mx) \cos(2ny)}{\sqrt{\beta_2^2 + [\tau^2 + (1-\tau^2) \cos^2 y] \cos^2 x}}, \quad (15)$$

where we introduced the integers m and n such that $2m = r_1 - r_2$, $2n = r_1 + r_2$. For $\tau = 0$, $\mathcal{R}_2(\mathbf{r})$ coincides with that of Ref.¹². On the other hand, in the case of quasi-one-dimensional zig-zag chains (i.e. $\tau = 1$), $\mathcal{R}_2(\mathbf{r})$ is nonzero only along direction $n = 0$, and in this direction it amounts to $\mathcal{R}_1(\mathbf{r})$ but with twice as large hopping as in the uniform one-dimensional case.

By expanding the square root in the denominator, the double integral of (15) can be related to a single integral involving a hypergeometric function ${}_3F_2$ ¹⁴:

$$\begin{aligned} \mathcal{R}_2(\mathbf{r}) &= \frac{(-1)^m (1-\tau^2)^{-n}}{2\pi} \frac{\Gamma(m+1/2)}{(2\beta_2)^{2m+1} \Gamma(n+1/2) \Gamma(m-n+1)} \\ &\times \int_{\tau^2}^1 dv v^{m-n} (1-v)^{n-1/2} (v-\tau^2)^{n-1/2} \\ &\times {}_3F_2\left(m+\frac{1}{2}, m+\frac{1}{2}, m+1; 2m+1, m-n+1; -\frac{v}{\beta_2^2}\right), \end{aligned} \quad (16)$$

for $m \geq n$, and

$$\begin{aligned} \mathcal{R}_2(\mathbf{r}) &= \frac{(-1)^n (1-\tau^2)^{-n}}{2\pi} \frac{\Gamma(2n+1)}{(2\beta_2)^{2n+1} \Gamma(n+m+1) \Gamma(n-m+1)} \\ &\times \int_{\tau^2}^1 dv (1-v)^{n-1/2} (v-\tau^2)^{n-1/2} \\ &\times {}_3F_2\left(n+\frac{1}{2}, n+\frac{1}{2}, n+1; n+m+1, n-m+1; -\frac{v}{\beta_2^2}\right), \end{aligned} \quad (17)$$

for $n \geq m$.

Using representations (16), (17), we have derived large-distance asymptotic behavior of $\mathcal{R}_2(\mathbf{r})$ in diagonal directions of a connected system ($\tau \neq 1$). Specifically, in direction $n = 0$ it follows from (16) that

$$\mathcal{R}_2(m, -m) \approx \frac{(-1)^m}{2\pi\beta_2} \frac{1}{\sqrt{1-\tau^2}} \frac{\exp(-2m/\xi)}{2m}, \quad (18)$$




FIG. 1: The inverse correlation length of D=2 system in direction $n = m$ against u , for different values of τ . Symbols represent the curves obtained from fitting numerically $\mathcal{R}(2m, 0)$. The continuous lines are derived from approximate formula (22), the upper one for $\tau = 0.25$, the lower one for $\tau = 0.05$. The dashed curve is derived from an exact formula for $\tau = 0$. In the inset we display the corresponding log-log plots.

$$\xi^{-1} = \ln(\sqrt{1+\beta_2^2} + \beta_2). \quad (19)$$

Apparently, in direction $n = 0$ the gap determines ξ . The above asymptotics differs only by the factor $(1-\tau^2)^{-1/2}$ from that in the uniform hopping case¹², hence it is continuous at $\tau = 0$. More interesting is a neighborhood of $\tau = 1$, where we would like to observe how our $D = 2$ system approaches the corresponding $D = 1$ system. In particular, an interesting question is how the power law r^γ , with $r = 2m$ and $\gamma = 1$ in (18), transforms into $\sqrt{2m}$ in (8), as $\tau \rightarrow 1$? Our numerical calculations suggest that γ behaves discontinuously: it grows to some value above 1, and then jumps to $1/2$ at $\tau = 1$.

Then, in direction $m = 0$ formula (17) gives

$$\mathcal{R}_2(n, n) \approx \frac{(-1)^n \exp(-2n/\xi)}{2\pi\beta_2} \frac{1}{2n}, \quad (20)$$

where ξ^{-1} is given by (9) with κ_1 replaced by κ_2 and κ_2 obtains from κ_1 by changing β_1 into β_2 . Similarly, we introduce δ_2 , $\delta_2^2 = \beta_2^2 + \tau^2$. While δ_2 and κ_2 lack the physical interpretation of δ_1 and κ_1 , the formal relation between them is fairly analogous to that between the latter parameters. Consequently, in direction $m = 0$ the gap does not determine ξ^{-1} .

In directions other than diagonal ones, we have not been able to establish the asymptotic behavior of the correlation function. But with the help of numerical calculations, we have obtained an approximate asymptotics

in the axial direction $m = n$. This asymptotics can be considered as an upper bound for the true one, and it approaches the true one as u grows and/or τ decreases. Specifically, after expanding the hypergeometric function of (16) (or (17)) in a series¹⁴, and carrying out the integration, we get a power series in $-1/\beta_2^2$ whose coefficient by power j depends on the hypergeometric function $F(-j, m+1/2; 2m+1; 1-\tau^2)$. One can verify numerically that if $\tau \ll 1$ and $j \ll m$, then this function is approximately equal to $F(-j, m+1/2; 2m+1; 1)(1+\tau^2)^j$. On substituting this approximation to the series and summing up the series, we arrive at an expression similar to formula (9) of Ref.¹², which in turn constitutes an approximation to $\mathcal{R}_2(2m, 0)$. The obtained approximation for the correlation function is the better the smaller is τ and/or the larger is u (see Fig.1). In particular, if $\tau = 0$, it becomes exact for any β_2 . On the basis of the results of Ref.¹², the approximate asymptotic behavior of $\mathcal{R}_2(2m, 0)$ has the form:

$$\mathcal{R}_2(2m, 0) \approx \frac{(-1)^m}{\pi \sqrt{2\beta_2}} \frac{1}{\sqrt[4]{1+\tau^2}} \frac{\exp(-2m/\xi)}{2m} \times \cos\left(m \arccos \frac{a-1}{a+1} - \frac{\pi}{4}\right), \quad (21)$$

where

$$\xi^{-1} = \ln \left[(\sqrt{a} + \sqrt{1+a}) \sqrt{\frac{1+\tau^2}{1-\tau^2}} \right], \quad a \equiv \frac{2\beta_2(\sqrt{1+\beta_2^2+\tau^2} + \beta_2)}{1+\tau^2}. \quad (22)$$

As can be seen in Fig.1, the above approximate ξ^{-1} is a lower bound for the numerically exact one. The asymptotic behaviors of the inverse correlation length (22), for

small and large gaps, read:

$$\xi^{-1} \approx \ln \sqrt{\frac{1+\tau^2}{1-\tau^2}} + \frac{\sqrt{2\beta_2}}{(1+\tau^2)^{1/4}} \left(1 + \frac{1}{6} \frac{\beta_2}{\sqrt{1+\tau^2}} + \dots\right),$$

$$\xi^{-1} \approx \ln \frac{4}{\sqrt{1-\tau^2}} + \ln \beta_2 + \frac{3}{16} \frac{\tau^2}{\beta_2^2} - \frac{35}{512} \frac{\tau^4}{\beta_2^4} + \dots, \quad (23)$$

respectively.

To summarize, continuing our studies of the decay rates of one-body reduced density matrices in insulators, by means of analytic and numerical methods, we have investigated a tight-binding model, where not only an external potential but also hoppings can be spatially modulated. The general result of our investigations, presented in this report and in Ref.¹², reads: the form of the large-distance decay is always a power law times an exponential, unless the hoppings and the external potential are simultaneously uniform. We have determined analytically the power in the power law and the correlation length for $D = 1$ system and in several directions for $D = 2$ system. In comparison to the uniform hopping case, the considered here system with modulated hoppings exhibits interesting novel features. Firstly, since the correlation length depends on two energy parameters (controlling the amplitudes of spatial modulation of the external potential and the hopping), in $D = 1$ case and in some directions of $D = 2$ case the gap does not determine the correlation length. Secondly, by adjusting the hopping-modulation amplitude suitably, the $D = 2$ -system can be decoupled into quasi-one-dimensional zig-zag chains, which has made possible investigating a crossover from $D = 2$ -decay rates to $D = 1$ ones.

T.K. is grateful to the Institute of Theoretical Physics of the University of Wrocław for kind hospitality and financial support.

* Electronic address: jjed@ift.uni.wroc.pl

† Electronic address: krokhn@icmp.lviv.ua

¹ M.C. Payne, M.P. Teter and D.C. Allan, T.A. Arias and J.D. Joannopoulos, Rev. Mod. Phys. **64**, 1045 (1992)

² S. Goedecker, Rev. Mod. Phys. **71**, 1085 (1999)

³ W. Kohn, Phys. Rev. Lett. **76**, 3168 (1996)

⁴ W. Kohn, Phys. Rev. **115**, 809 (1959)

⁵ R. Baer and M. Head-Gordon, Phys. Rev. Lett. **79**, 3962 (1997)

⁶ S. Goedecker, Phys. Rev. **B 58**, 3501 (1998)

⁷ S. Ismail-Beigi and T. A. Arias, Phys. Rev. Lett. **82**, 2127 (1999)

⁸ L. He and D. Vanderbilt, Phys. Rev. Lett. **86**, 5341 (2001)

⁹ X. Zhang and D. A. Drabold, Phys. Rev. **B 63**, 233109 (2001)

¹⁰ S. N. Taraskin, D. A. Drabold, and S. R. Elliott, Phys. Rev. Lett. **88**, 196405 (2002)

¹¹ S. N. Taraskin, P. A. Fry, X. Zhang, D. A. Drabold, and S. R. Elliott, Phys. Rev. **B 66**, 233101 (2002)

¹² J. Jędrzejewski and T. Krokhmal'skii, Phys. Rev. **B 70**, 153102 (2004)

¹³ J. Des Cloizeaux, Phys. Rev. **135**, A698 (1964)

¹⁴ H. Bateman and A. Erdelyi, *Higher transcendental functions*, vol. 1, McGraw-Hill, New York 1953